


**AMENDMENTS TO THE CLAIMS**

The following is a marked-up version of the claims with the language that is underlined ("\_\_\_") being added and the language that contains strikethrough ("~~—~~") being deleted:


1. (Currently Amended) A compound selected from the group consisting of: a compound of Formula I, a pharmaceutically acceptable salt thereof, and a stereoisomer thereof:



**Formula I**

wherein n is an integer;

wherein  $R_1$  and  $R_2$  are independently selected from the group consisting of:




wherein R<sub>3</sub> is selected from the group consisting of: H; alkyl; alkenyl; alkynyl; halogen; aryl; heteroaryl containing N, O, or S; NZ<sub>1</sub>Z<sub>2</sub>, wherein Z<sub>1</sub> and Z<sub>2</sub> are independently selected from the group consisting of: H and alkyl; and (CO)Y<sub>1</sub> wherein Y<sub>1</sub> is selected from the group consisting of: H, alkyl, alkenyl, alkynyl, aryl, heteroaryl containing N, O, or S; and

wherein R<sub>4</sub> is selected from the group consisting of: H and alkyl.


2. (Original) The compound of claim 1, wherein n is 1.
3. (Original) The compound of claim 1, wherein the aryl and heteroaryl are substituted with at least one of a halogen, an alkyl, an alkenyl, and an alkynyl.

4. (Currently Amended) The compound of claim 1, wherein the compound is selected from the group consisting of: a compound of Formula II, a pharmaceutically acceptable salt thereof, and a stereoisomer thereof:




Formula II

5. (Currently Amended) The compound of claim 1, wherein the compound is selected from the group consisting of: a compound of Formula III, a pharmaceutically acceptable salt thereof, and a stereoisomer thereof:



Formula III

6. (Currently Amended) The compound of claim 1, wherein the compound is selected from the group consisting of: a compound of Formula IV, a pharmaceutically acceptable salt thereof and a stereoisomer thereof:



Formula IV

7. (Original) The compound of claim 1, wherein the pharmaceutically acceptable salt is derived from an inorganic acid or an organic acid, wherein the inorganic acid is selected from the group consisting of hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric acids; and the organic acid is selected from the group consisting of acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and trifluoroacetic acids.


8. (Original) The compound of claim 7, wherein the pharmaceutically acceptable salt is derived from hydrochloric acid.

9. – 10. (Canceled)


11. (Original) A pharmaceutical composition comprising the compound of claim 1 or mixtures thereof, and a pharmaceutically acceptable carrier.

12. (Original) A method for making the compound of claim 1, comprising reacting  
with

in the presence of a palladium coupling catalyst for promoting carbon-nitrogen bond-forming cross-coupling.



13. (Original) The method of claim 12, wherein the palladium coupling catalyst is one of



$\text{Pd}(\text{OAc})_2$  with BINAP and  $\text{Pd}_2(\text{dba})_3$  with DPPF.

14. (Original) The method of claim 12, wherein the  $\text{R}_1$  and/or  $\text{R}_2$  groups contain protecting groups and the method further comprises a deprotection step.

15. (Currently Amended) A method of treating a tyrosine kinase-dependent cancer in a mammal in need of such treatment, wherein the cancer is selected from the group of: cancers consisting of breast, colon, ovarian, prostate, chronic myeloid leukemia (CML), leukemia, melanoma, CNS, and lung, comprising administering to the mammal a therapeutically effective amount of the compound of claim 1 or mixtures thereof.

16. (Original) The method of claim 15, wherein the therapeutically effective amount is between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day.
17. (Original) The method of claim 16, wherein the therapeutically effective amount is between about 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.
18. (Canceled)
19. (Currently Amended) The method of claim 15 further comprising a therapy selected from the group consisting of: radiation therapy and chemotherapy.
20. (Original) The method of claim 19, wherein the therapeutically effective amount is between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day.
21. (Original) The method of claim 20, wherein the therapeutically effective amount is between about 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.
22. (Canceled)
23. (Previously Presented) The method of claim ~~18~~ 15, wherein the cancer is chronic myeloid leukemia (CML).
24. (Original) A process for making a pharmaceutical composition which comprises combining a compound of claim 1 or mixtures thereof, with a pharmaceutically acceptable carrier.
25. (Currently Amended) A composition comprising a compound of claim 1 or mixtures thereof, and a compound selected from the group consisting of: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic agent, an anti-proliferative agent, a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of

platelet derived growth factor, an MMP inhibitor, an integrin blocker, interferon- $\alpha$ , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, and troponin-1, tamoxifen and raloxifene.

26.-35. (Canceled)